arXiv:1304.3070v1 [math.GT] 10 Apr 2013

LESCOP'S INVARIANT AND GAUGE THEORY

PRAYAT POUDEL

Abstract. Taubes proved that the Casson invariant of an integral homology 3-sphere equals half the Euler characteristic of its instanton Floer homology. We extend this result to all closed oriented 3-manifolds with positive first Betti number by establishing a similar relationship between the Lescop invariant of the manifold and its instanton Floer homology. The proof uses surgery techniques.

1. Introduction

The Lescop invariant $\lambda_L(M)$ is a rational valued invariant of closed oriented 3-manifolds M defined by Lescop [9] via a combinatorial formula as a generalization of the Casson invariant [1]. The Casson invariant, while only defined for integral homology 3-spheres, has a very useful gauge theoretic interpretation, due to Taubes [14], as half the Euler characteristic of the instanton Floer homology [5]. We provide a similar interpretation of the Lescop invariant for all 3-manifolds with the positive first Betti number using a version of instanton Floer homology based on admissible bundles. In fact, our formula matches the one conjectured in the physics literature, where the Lescop invariant arises as a partition function of the Donaldson-Witten theory of a 4-manifold of the form $S^1 \times M$, see Mariño–Moore [10].

Theorem. *Let M be a closed oriented connected 3-manifold with $b_1(M) \geq 1$, and let $\lambda_L(M)$ be its Lescop invariant. Then there exists an admissible bundle P over M such that*

$$\lambda_L(M) = -\frac{1}{2}\,\chi(I_*\,(M,P)) - \frac{1}{12}\,|\operatorname{Tor}(H_1(M))|, \quad \text{if } b_1(M) = 1, \text{ and}$$

$$\lambda_L(M) = -\frac{1}{2}\,(-1)^{b_1(M)} \cdot |\operatorname{Tor}(H_1(M))| \cdot \chi(I_*\,(M,P)), \quad \text{if } b_1(M) \geq 2,$$

2010 *Mathematics Subject Classification.* 57M27, 57R58.

Key words and phrases. Casson invariant, Lescop invariant, Floer homology.

where $\chi(I_*(M,P))$ *stands for the Euler characteristic of the instanton Floer homology of the pair* (M,P)*, see Section 2.*

In addition, if $H_1(M)$ has no 2-torsion, we show that $\chi(I_*(M,P))$ is independent of the choice of admissible bundle P hence the above formulas hold for *any* admissible bundle P. We conjecture that the same is true for any M with $b_1(M) \geq 1$, with or without 2-torsion in its homology.

Still lacking is a gauge theoretic interpretation of the Lescop invariant for rational homology 3-spheres with non-trivial torsion because there is no satisfactory definition of instanton Floer homology for such manifolds.

Our proof will proceed by induction on the first Betti number $b_1(M)$ of the manifold and will use the Floer exact triangle [6]. In case M has no torsion in homology, we will start the induction at $b_1(M) = 0$ and use Taubes' theorem [14]. In the presence of torsion, due to the aforementioned problem with defining instanton Floer homology for rational homology spheres, we will start at $b_1(M) = 1$ and use an extension of Taubes' theorem due to Masataka [11].

Acknowledgments. I am immensely grateful to my Ph.D. advisor Dr. Nikolai Saveliev for his guidance and support. I am also thankful to Dr. Ken Baker and Dr. Kim Frøyshov for their helpful suggestions.

2. INSTANTON FLOER HOMOLOGY

Let M be a closed oriented connected 3-manifold and $P \to M$ a $U(2)$-bundle such that one of the following conditions holds:

1. M is an integral homology sphere and P is a trivial $SU(2)$-bundle, or
2. $b_1(M) \geq 1$ and P is a $U(2)$-bundle whose first Chern class $c_1(P)$ has an odd pairing with some integral homology class in $H_2(M)$. Note that the second Stiefel-Whitney class $w_2(\mathrm{ad}(P)) \in H^2(M;\mathbb{Z}_2)$ of the associated $SO(3)$-bundle $\mathrm{ad}(P)$ is then not zero as a map $H_2(M) \to \mathbb{Z}_2$.

Both the bundle P and its adjoint bundle $\mathrm{ad}(P)$ will be referred to as admissible bundles.

Given an admissible bundle P, consider the space $\mathcal{C}$ of $SO(3)$-connections in $\mathrm{ad}(P)$. This space is acted upon by the group $\mathcal{G}$ of determinant one gauge transformations of P. The instanton Floer homology $I_*(M, P)$ is the Floer homology arising from the Chern-Simons function of the space $\mathcal{C}/\mathcal{G}$, see Donaldson [4].

The groups $I_*(M, P)$ depend on the choice of $\mathrm{ad}(P)$ but not P. They have an absolute $\mathbb{Z}_2$-grading defined as in Section 5.6 of [4]. These groups also admit a relative $\mathbb{Z}_8$-grading, which becomes an absolute $\mathbb{Z}_8$-grading if M is an integral homology sphere. Note that our setup is consistent with that of Kronheimer and Mrowka [8], Section 7.1. Using their notations, $I_*(M, P) = I_*(M)_w$, where $w = \det(P)$ is the determinant bundle of P.

Two main ingredients that go into our calculation of the Euler characteristic of $I_*(M, P)$ are as follows. The first one is a special surgery presentation of M as in [9], Lemma 5.1.1.

Lemma 2.1. *Let M be a closed connected oriented 3-manifold and $b_1(M) = n \geq 0$. Then there exists a rational homology sphere Σ and an algebraically split n-component link $\mathcal{L} \subset \Sigma$ such that $M = \Sigma + 0 \cdot \mathcal{L}$, each component of $\mathcal{L}$ is null-homologous in Σ, and $|H_1(\Sigma)| = |\operatorname{Tor}(H_1(M))|$.*

We denote by $\Sigma + 0 \cdot \mathcal{L}$ the manifold obtained by 0-framed surgery on $\mathcal{L}$. Similarly, $\Sigma \pm \ell$ will denote the result of (± 1)-surgery on knot ℓ.

The second ingredient is a long exact sequence known as the Floer exact triangle, see [2]. Let Σ be a rational homology sphere and $\mathcal{L} = \ell_1 \cup \ell_2 \ldots \cup \ell_n$ be an algebraically split link in Σ. If $n = 1$, we will require that Σ be an integral homology sphere. Then we have the following Floer exact triangle:

$$\begin{tikzcd}
& I_*(\Sigma + 0 \cdot \ell_1 + \ldots + 0 \cdot \ell_n) \arrow[dl, "-1"'] & \\
I_*\left(\Sigma + 0 \cdot \ell_1 + \ldots + 0 \cdot \ell_{n-1}\right) \arrow[rr] & & I_*\left((\Sigma + 0 \cdot \ell_1 + \ldots 0 \cdot \ell_{n-1}) - \ell_n\right) \arrow[ul]
\end{tikzcd}$$

The admissible bundles P, which are omitted from the notations, have $c_1(P)$ pair non-trivially with the natural homology classes obtained by capping off a Seifert surface of ℓ_j by a meridional disk of the surgery. In addition, the three admissible bundles that show up in the Floer exact triangle match when restricted to the exterior of the link $\ell_1 \cup \cdots \cup \ell_n$ in Σ. If $n = 1$, the bundles P over Σ and $\Sigma - \ell_1$ are trivial $SU(2)$-bundles.

It should be pointed out that we will not use the complete strength of the Floer exact triangle; all we will derive from it is the following relation on Euler characteristics:

$$\chi(I_*(\Sigma + 0\cdot\ell_1 + 0\cdot\ell_2 + \ldots + 0\cdot\ell_n)) =$$
$$\chi(I_*(\Sigma + 0\cdot\ell_1 + \ldots + 0\cdot\ell_{n-1}) - \ell_n) - \chi(I_*(\Sigma + 0\cdot\ell_1 + \ldots + 0\cdot\ell_{n-1})).$$

The obvious observation that

$$\begin{aligned} &b_1((\Sigma + 0\cdot\ell_1 + \ldots + 0\cdot\ell_{n-1}) - \ell_n)) = b_1(\Sigma + 0\cdot\ell_1 + \ldots + 0\cdot\ell_{n-1}) \\ &\quad = b_1(\Sigma + 0\cdot\ell_1 + 0\cdot\ell_2 + \ldots 0\cdot\ell_n) - 1 \end{aligned}$$

allows us to proceed via induction on $b_1(M)$.

3. CASE OF $b_1(M) = 0$

If M is an integral homology sphere, then $\lambda_L(M) = \lambda(M)$, which is the Casson invariant of M, see Section 1.5 of [9]. On the Floer homology side, we work with the trivial $SU(2)$-bundle P over M, and denote the instanton Floer homology by $I_*(M,P)$. According to Taubes [14], we have $\chi(I_*(M,P)) = 2\cdot\lambda(M)$. Therefore, $\chi(I_*(M,P)) = 2\cdot\lambda_L(M)$.

4. CASE OF $b_1(M) = 1$

Let $k \subset \Sigma$ be a null-homologous knot in a rational homology sphere Σ. Choose a Seifert surface F of k, and denote by V its Seifert matrix with respect to a basis of $H_1(F)$. The Laurent polynomial

$$\Delta_{k\subset\Sigma}(t) = |H_1(\Sigma)| \cdot \det(t^{1/2}\,V - t^{-1/2}\,V^\top)$$

is called the Alexander polynomial of $k \subset \Sigma$. Note that $\Delta_{k\subset\Sigma}(t) = \Delta_{k\subset\Sigma}(-t)$ and $\Delta_{k\subset\Sigma}(1) = |H_1(\Sigma)| > 0$.

Given a closed oriented connected 3-manifold M with $b_1(M) = 1$, according to Lemma 2.1, there exists a null-homologous knot $\ell_1 \subset \Sigma$ in a rational homology sphere Σ such that $M = \Sigma + 0\cdot\ell_1$. The Lescop invariant of M is then equal to

$$\lambda_L(M) = \frac{1}{2}\Delta''_{\ell_1\subset\Sigma}(1) - \frac{1}{12}\,|\operatorname{Tor}(H_1(M))|, \tag{1}$$

see [9], Section 1.5. It is independent of the choice of surgery presentation of M.

In the special case when Σ is an integral homology sphere, it follows from the Floer exact triangle and Casson's surgery formula [1] that

$$\begin{aligned}\chi(I_*(M,P)) &= \chi(I_*(\Sigma-\ell_1)) - \chi(I_*(\Sigma)) = 2\lambda(\Sigma-\ell_1) - 2\lambda(\Sigma)\\ &= 2\left(-\frac{1}{2}\Delta''_{\ell_1\subset\Sigma}(1)\right) = -\Delta''_{\ell_1\subset\Sigma}(1).\end{aligned}$$

Therefore,

$$\chi(I_*(M,P)) = -2\lambda_L(M) - \frac{1}{6}\left|\mathrm{Tor}(H_1(M))\right| \tag{2}$$

as claimed, for the unique admissible $SU(2)$-bundle P over M. The general case is handled similarly using the following result.

Proposition 4.1. *Let* $M = \Sigma + 0\cdot\ell_1$*, where* ℓ_1 *is a null-homologous knot in a rational homology sphere* Σ*. Then* $\chi(I_*(M,P)) = -\Delta''_{\ell_1\subset\Sigma}(1)$ *for any admissible bundle* $\mathrm{ad}(P)$ *over* M *whose restriction to the exterior of* $\ell_1\subset\Sigma$ *is trivial.*

Proof. If $H_1(\Sigma)$ has non-trivial torsion, the starting point for our calculation will be the result from [11] which, with our normalization, reads

$$\frac{1}{2}\chi(I_*(M,P)) = -\lambda_L(\Sigma+\ell_1) + \lambda_L(\Sigma).$$

We wish to identify the right hand side of this equation with $-\frac{1}{2}\Delta''_{\ell_1\subset\Sigma}(1)$. By Lescop [9],

$$\lambda_L(\Sigma+\ell_1) = \frac{|H_1(\Sigma+\ell_1)|}{|H_1(\Sigma)|}\cdot\lambda_L(\Sigma) + \mathbb{F}_\Sigma(\ell_1),$$

where $\mathbb{F}_\Sigma(\ell_1)$ is defined by equation 1.4.8 in [9]. Since $|H_1(\Sigma+\ell_1)| = |H_1(\Sigma)|$, we conclude that $\lambda_L(\Sigma+\ell_1) = \lambda_L(\Sigma) + \mathbb{F}_\Sigma(\ell_1)$ and therefore

$$\frac{1}{2}\chi(I_*(M,P)) = -\mathbb{F}_\Sigma(\ell_1).$$

A straightforward calculation of $\mathbb{F}_\Sigma(\ell_1)$ shows that

$$\mathbb{F}_\Sigma(\ell_1) = \frac{1}{2}\Delta''_{\ell_1\subset\Sigma}(1),$$

which leads to the desired formula. Therefore, for our choice of admissible bundle P over M,

$$\chi(I_*(M,P)) = -2\lambda_L(M) - \frac{1}{6}\left|\mathrm{Tor}(H_1(M))\right|.$$

□

Remark. The exact sequence of the universal coefficient theorem,

$$0 \longrightarrow \operatorname{Ext}(H_1(M), \mathbb{Z}_2) \longrightarrow H^2(M; \mathbb{Z}_2) \longrightarrow \operatorname{Hom}(H_2(M), \mathbb{Z}_2) \longrightarrow 0,$$

tells us that the ambiguity in choosing an admissible bundle $\operatorname{ad}(P)$ over $M = \Sigma + 0 \cdot \ell_1$ resides in the group $\operatorname{Ext}(H_1(M), \mathbb{Z}_2)$. If $H_1(M)$ has no 2-torsion, the latter group vanishes, implying that there is a unique admissible bundle $\operatorname{ad}(P)$ over M.

5. CASE OF $b_1(M) = 2$

Let $\mathcal{L} = \ell_1 \cup \ell_2$ be an oriented two-component link in a rational homology sphere Σ such that ℓ_1, ℓ_2 are null-homologous in Σ and $\ell k_\Sigma(\ell_1, \ell_2) = 0$. There exist Seifert surfaces F_1 and F_2 of ℓ_1 and ℓ_2, respectively, such that $F_1 \cap \ell_2 = \emptyset$ and $F_2 \cap \ell_1 = \emptyset$. If the surfaces intersect, they may be assumed to intersect in a circle c, see [3]. The self linking number of c with respect to either F_1 or F_2 is called the Sato-Levine invariant and is denoted by $s(\ell_1 \cup \ell_2 \subset \Sigma)$. To be precise, $s(\ell_1 \cup \ell_2 \subset \Sigma) = \ell k_\Sigma(c, c^+)$, where c^+ is a positive push-off of c with respect to either F_1 or F_2. If the surfaces don't intersect then $s(\ell_1 \cup \ell_2 \subset \Sigma)$ =0.

Given a closed oriented connected 3-manifold M with $b_1(M) = 2$, according to Lemma 2.1, there exists an algebraically split link $\ell_1 \cup \ell_2 \subset \Sigma$ in a rational homology sphere Σ such that $M = \Sigma + 0 \cdot \ell_1 + 0 \cdot \ell_2$. According to Lescop [9], Section 5.1,

$$\lambda_L(M) = -|\operatorname{Tor}(H_1(M))| \cdot s(\ell_1 \cup \ell_2 \subset \Sigma); \tag{3}$$

it is independent of the choice of surgery presentation of M. We will first handle the case when Σ is an integral homology sphere.

Proposition 5.1. *Let $M = \Sigma + 0 \cdot \ell_1 + 0 \cdot \ell_2$, where $\ell_1 \cup \ell_2$ is a link in an integral homology sphere Σ such that $\ell k_\Sigma(\ell_1, \ell_2) = 0$. Then, for any admissible bundle* $\operatorname{ad}(P)$ *over M,*

$$\chi(I_*(M, P)) = 2 \cdot \lambda_L(M).$$

Proof. Let P be a $U(2)$-bundle over M such that $w_2(\operatorname{ad}(P))$ evaluates non-trivially on both homology classes obtained by capping off Seifert surfaces of ℓ_1 and ℓ_2 by meridional disks of the surgery. We will indicate this by writing

$w_2(\mathrm{ad}(P)) = (1,1)$ in the natural basis of $\mathrm{Hom}(H_2(M),\mathbb{Z}_2)$. It follows from the Floer exact triangle that

$$\begin{aligned}\chi(I_*\,(M,P)) &= \chi(I_*\,((\Sigma + 0\cdot\ell_1) - \ell_2)) - \chi(I_*\,(\Sigma + 0\cdot\ell_1))\\ &= -\Delta''_{\ell_1\subset\Sigma-\ell_2}(1) + \Delta''_{\ell_1\subset\Sigma}(1),\end{aligned}$$

hence the calculation boils down to computing Alexander polynomials of $\ell_1 \subset \Sigma$ and $\ell_1 \subset \Sigma - \ell_2$.

According to Hoste [7], there exist Seifert surfaces F_1 and F_2 of ℓ_1 and ℓ_2, respectively, such that $F_1 \cap F_2$ is either empty or a single ribbon intersection, and furthermore, $F_1 \cap F_2$ does not separate F_1 or F_2.

If $F_1 \cap F_2$ is empty then $\ell_1 \cup \ell_2$ is a boundary link and $s(\ell_1 \cup \ell_2 \subset \Sigma) = 0$. Moreover, (-1)-surgery along ℓ_2 leaves F_1 unaffected, therefore, $\Delta_{\ell_1\subset\Sigma}(t) = \Delta_{\ell_1\subset\Sigma-\ell_2}(t)$ and $\chi(I_*\,(M,P)) = 0$ as desired.



Figure 1.



Figure 2.

If $F_1 \cap F_2$ is non-empty then it is a single ribbon intersection as in Figure 1. The intersection $F_1 \cap \partial F_2 \subset \ell_2$ consists of two points that separate ℓ_2 into two arcs, ℓ_2' and ℓ_2''. Stabilize surface F_1 by adding a tube with core ℓ_2' and call this new modified surface F_1', see Figure 2. If F_1 had genus g, the genus of F_1' will be $g+1$.

The intersection $F_1' \cap F_2$ is a closed loop c which represents a primitive homology class in $H_1(F_1')$. Complete c to a basis $\{e_1, c, e_3, \ldots, e_{2g+2}\}$ of $H_1(F_1')$, where e_1 is a meridional curve of ℓ_2 and $\{e_3, \ldots, e_{2g+2}\}$ is a basis for $H_1(F_1)$. Using general position argument we will assume that $e_m \cap F_2 =$

$\emptyset$ and hence $\ell k_\Sigma(e_m, \ell_2) = 0$ for $m \geq 3$. In addition, it is obvious that $\ell k_\Sigma(e_1, \ell_2) = \pm 1$, and that $\ell k_\Sigma(c, \ell_2) = \ell k_\Sigma(c^+, \ell_2) = 0$, where c^+ is a positive push-off of c with respect to F_2. To summarize, we have the matrix

$$E = \begin{bmatrix} \ell k_\Sigma(e_1, \ell_2) \\ \ell k_\Sigma(c, \ell_2) \\ \vdots \\ \ell k_\Sigma(e_{2g+2}, \ell_2) \end{bmatrix} = \begin{bmatrix} \pm 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

This matrix accounts for the difference in the Seifert matrices of ℓ_1 when viewed as a knot in Σ and $\Sigma - \ell_2$. To be precise, according to Hoste [7],

$$V(\ell_1 \subset \Sigma - \ell_2) = V(\ell_1 \subset \Sigma) + E \cdot E^\top. \tag{4}$$

We would like next to determine the Seifert matrix $V(\ell_1 \subset \Sigma)$ with respect to the basis $\{e_1, c, e_3, \ldots e_{2g+2}\}$ in $H_1(F_1')$. To begin, $\ell k_\Sigma(e_1, e_1^+) = 0$ since the push off e_1^+ is disjoint from the meridional disk D of e_1. For $m \geq 3$, the curve e_m lies on the surface F_1. Therefore, its push-off e_m^+ can be isotoped to make it disjoint from D so $\ell k_\Sigma(e_1, e_m^+) = 0$. Similarly $\ell k_\Sigma(e_m, e_1^+) = 0$ for $m \geq 3$. Next, $\ell k_\Sigma(c, e_1^+) = \pm 1$ and since c^+ does not intersect the meridional disk bounded by e_1, $\ell k_\Sigma(e_1, c^+) = 0$. To finish the calculation, observe that $V(\ell_1 \subset \Sigma) - V(\ell_1 \subset \Sigma)^\top = -I$, where $I : H_1(F_1') \times H_1(F_1') \to \mathbb{Z}$ denotes the intersection form given by $I(v, w) = v \cdot w$. By general position argument, $c \cdot e_m = 0$ hence $\ell k_\Sigma(c^+, e_m) - \ell k_\Sigma(c, e_m^+) = 0$. For $m \geq 3$, denote $\ell k_\Sigma(c^+, e_m) = \ell k_\Sigma(c, e_m^+) = a_{m-2}$. Finally, $\ell k_\Sigma(c, c^+) = s$ by the definition of $s = s(\ell_1 \cup \ell_2 \subset \Sigma)$. Therefore, we obtain the matrix

$$V(\ell_1 \subset \Sigma) = \left[\begin{array}{cc|ccc} 0 & 0 & 0 & \cdots & 0 \\ \pm 1 & s & a_1 & \cdots & a_{2g} \\ \hline 0 & a_1 & & & \\ \vdots & \vdots & & W & \\ 0 & a_{2g} & & & \end{array}\right]$$

where W is the Seifert matrix of ℓ_1 with respect to the basis $e_3, \ldots e_{2g} \in H_1(F_1)$. Using Hoste's formula (4), we obtain

$$V(\ell_1 \subset \Sigma - \ell_2) = \left[\begin{array}{cc|ccc} 1 & 0 & 0 & \cdots & 0 \\ \pm 1 & s & a_1 & \cdots & a_{2g} \\ \hline 0 & a_1 & & & \\ \vdots & \vdots & & & \\ & & & W & \\ 0 & a_{2g} & & & \end{array}\right]$$

Now we are ready to compute the Alexander polynomials. Denote $V(\ell_1 \subset \Sigma - \ell_2)$ by V and let $z = t^{1/2} - t^{-1/2}$. Then

$$t^{1/2}V - t^{-1/2}V^{\top} = \left[\begin{array}{cc|ccc} z & \mp t^{-1/2} & 0 & \cdots & 0 \\ \pm t^{1/2} & sz & a_1 z & \cdots & a_{2g} z \\ \hline 0 & a_1 z & & & \\ \vdots & \vdots & & t^{1/2}W - t^{-1/2}W^{\top} & \\ 0 & a_{2g} z & & & \end{array}\right]$$

Hence $\det(t^{1/2}V - t^{-1/2}V^{\top}) =$

$$= z \left|\begin{array}{c|ccc} sz & a_1 z & \cdots & a_{2g} z \\ \hline a_1 z & & & \\ \vdots & & & \\ & & t^{1/2}W - t^{-1/2}W^{\top} & \\ a_{2g} z & & & \end{array}\right| \mp t^{-1/2} \left|\begin{array}{c|ccc} \mp t^{-\frac{1}{2}} & 0 & \cdots & 0 \\ \hline a_1 z & & & \\ \vdots & & & \\ & & t^{1/2}W - t^{-1/2}W^{\top} & \\ a_{2g} z & & & \end{array}\right|$$

$$= z^3 \left|\begin{array}{c|ccc} sz^{-1} & a_1 & \cdots & a_{2g} \\ \hline a_1 & & & \\ \vdots & & & \\ & & t^{1/2}W - t^{-1/2}W^{\top} & \\ a_{2g} & & & \end{array}\right| + \det(t^{1/2}W - t^{-1/2}W^{\top})$$

$$= s\, z^2 \cdot \det(t^{1/2}W - t^{-1/2}W^{\top}) + z^3 \cdot f(t) + \Delta_{\ell_1 \subset \Sigma}(t)$$

for some function $f(t)$, which is a polynomial in $t^{1/2}$ and $t^{-1/2}$. Therefore,

$$\Delta_{\ell_1 \subset \Sigma - \ell_2}(t) = sz^2 \Delta_{\ell_1 \subset \Sigma}(t) + z^3 f(t) + \Delta_{\ell_1 \subset \Sigma}(t).$$

Next we differentiate twice and set $t = 1$. An easy calculation taking into account that $z(1) = 0$ and $z'(1) = 1$ leads to the formula

$$\Delta''_{\ell_1 \subset \Sigma - \ell_2}(1) = 2s \cdot \Delta_{\ell_1 \subset \Sigma}(1) + \Delta''_{\ell_1 \subset \Sigma}(1) = 2s + \Delta''_{\ell_1 \subset \Sigma}(1).$$

Therefore,

$$\chi(I_*\,(M, P)) = -2s(\ell_1 \cup \ell_2 \subset \Sigma) = 2 \cdot \lambda_L(M).$$

The above calculation holds for a specific bundle which has $w_2(\mathrm{ad}(P)) = (1, 1)$. We will now prove the result for admissible bundles with $w_2(\mathrm{ad}(P)) = (0, 1)$ and $w_2(\mathrm{ad}(P)) = (1, 0)$ using the fact that $\lambda_L(M) = -|\operatorname{Tor}(H_1(M))| \cdot s(\ell_1 \cup \ell_2 \subset \Sigma)$ is an invariant of the manifold M and therefore is independent of the surgery presentation.

Without loss of generality, let us assume that $w_2(\mathrm{ad}(P)) = (1, 0)$. After sliding ℓ_1 over ℓ_2, we will obtain a new surgery presentation for M, namely, $M = \Sigma + 0 \cdot \ell_1 + 0 \cdot \ell_\sharp$, where $\ell_\sharp$ is the new knot obtained by sliding ℓ_1 over ℓ_2. Note that $\ell_\sharp$ bounds a Seifert surface which is a band sum of F_1 and F_2 and also that $\ell k_\Sigma(\ell_1, \ell_\sharp) = 0$. In the new basis, $w_2(\mathrm{ad}(P)) = (1, 1)$, hence $\chi(I_*\,(M, P)) = s(\ell_1 \cup \ell_\sharp \subset \Sigma)$ by the above argument. The independence of surgery presentation then implies that $s(\ell_1 \cup \ell_\sharp \subset \Sigma) = s(\ell_1 \cup \ell_2 \subset \Sigma)$ and therefore $\chi(I_*\,(M, P))$ is independent of the choice of admissible bundle. $\square$

Before we move on to the case when $H_1(M)$ has non-trivial torsion, we will establish a fact which we will need for the calculations. Recall that if Σ is a rational homology sphere and k is an arbitrary knot in Σ, then there exists an integer q such that $q \cdot k$ represents a zero in $H_1(\Sigma)$. For any other knot ℓ in Σ, the linking number $\ell k_\Sigma(q \cdot k, \ell)$ is defined as the intersection number of a Seifert surface of $q \cdot k$ with ℓ, and one further defines

$$\ell k_\Sigma(k, \ell) = \frac{1}{q} \cdot \ell k_\Sigma(q \cdot k, \ell) \ \in \ \mathbb{Q}.$$

Lemma 5.2. *Let Σ be a rational homology sphere, $\ell_2 \subset \Sigma$ a null-homologous knot and k_1, k_2 knots in $\Sigma \smallsetminus N(\ell_2)$. Then,*

$$\ell k_{\Sigma - \ell_2}(k_1, k_2) = \ell k_\Sigma(k_1, k_2) + \ell k_\Sigma(k_1, \ell_2) \cdot \ell k_\Sigma(k_2, \ell_2). \tag{5}$$

Proof. If k_1, k_2 are null-homologous in $\Sigma \smallsetminus N(\ell_2)$, the proof proceeds exactly as in [7, Lemma 1.2]. If k_1 and k_2 are arbitrary knots in $\Sigma \smallsetminus N(\ell_2)$, there exist non-zero integers q_1, q_2 such that $q_1 \cdot k_1$ and $q_2 \cdot k_2$ are null-homologous in $\Sigma \smallsetminus N(\ell_2)$ and therefore in $\Sigma - \ell_2$. Then

$$\ell k_{\Sigma-\ell_2}(q_1 \cdot k_1, q_2 \cdot k_2) = \ell k_\Sigma(q_1 \cdot k_1, q_2 \cdot k_2) + \ell k_\Sigma(q_1 \cdot k_1, \ell_2) \cdot \ell k_\Sigma(q_2 \cdot k_2, \ell_2)$$

and the result follows by dividing both sides by $q_1 \cdot q_2$. □

Proposition 5.3. *Let* $M = \Sigma + 0 \cdot \ell_1 + 0 \cdot \ell_2$*, where* Σ *is a rational homology sphere and* $\ell_1, \ell_2 \subset \Sigma$ *are null-homologous knots such that* $\ell k_\Sigma(\ell_1, \ell_2) = 0$*. Then*

$$\chi(I_*(M, P)) = \frac{2 \cdot \lambda_L(M)}{|\operatorname{Tor}(H_1(M))|}$$

for any admissible bundle $\operatorname{ad}(P)$ *over* M *whose restriction to the exterior of* $\ell_1 \cup \ell_2 \subset \Sigma$ *is trivial.*

Proof. Let P be a $U(2)$-bundle over M such that $w_2(\operatorname{ad}(P))$ evaluates non-trivially on both homology classes obtained by capping off Seifert surfaces of ℓ_1 and ℓ_2 by meridional disks of the surgery. By the Floer exact triangle,

$$\begin{aligned}\chi(I_*(\Sigma + 0 \cdot \ell_1 + 0 \cdot \ell_2)) &= \chi(I_*((\Sigma + 0 \cdot \ell_1) - \ell_2)) - \chi(I_*(\Sigma + 0 \cdot \ell_1)) \\ &= -\Delta''_{\ell_1 \subset \Sigma - \ell_2}(1) + \Delta''_{\ell_1 \subset \Sigma}(1).\end{aligned}$$

We proceed as in the proof of Proposition 5.1 and choose Seifert surfaces F_1 and F_2 which are either disjoint or intersect in a single ribbon, and construct Seifert surfaces F_1' and F_2 which intersect in a circle c. Complete c to a basis $\{e_1, c, e_3, ..., e_{2g+2}\}$ of $H_1(F_1')$, where e_1 is a meridional curve of ℓ_2 and $\{e_3, ..., e_{2g+2}\}$ is a basis for $H_1(F_1)$. Again,

$$E = \begin{bmatrix} \ell k_\Sigma(e_1, \ell_2) \\ \ell k_\Sigma(c, \ell_2) \\ \vdots \\ \ell k_\Sigma(e_{2g+2}, \ell_2) \end{bmatrix} = \begin{bmatrix} \pm 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

We would like to determine the Seifert matrix $V(\ell_1 \subset \Sigma)$ with respect to the basis $\{e_1, c, e_3, ..., e_{2g+2}\}$ of $H_1(F_1')$. As e_1 bounds a meridional disk D, it is null-homologous in Σ. Since e_1^+ is disjoint from D, we have $\ell k_\Sigma(e_1, e_1^+) = 0$. As in the torsion free case, $\ell k_\Sigma(c, e_1^+) = 0$ and $\ell k_\Sigma(e_1, c^+) = \pm 1$. For $m \geq 3$, the disk D can be picked to be disjoint from e_m^+ hence $\ell k_\Sigma(e_1, e_1^+) = 0$ and similarly $\ell k_\Sigma(e_m, e_1^+) = 0$. By definition, $\ell k_\Sigma(c, c^+) = s$. Finally,

$\ell k_\Sigma(c^+, e_m) = \ell k_\Sigma(c, e_m^+)$ using the fact that $V(\ell_1 \subset \Sigma) - V(\ell_1 \subset \Sigma)^\top = -I$, where I is the intersection form on $H_1(F_1^{'}; \mathbb{Q})$. For $m \geq 3$ denote $\ell k_\Sigma(c^+, e_m) = \ell k_\Sigma(c, e_m^+) = a_{m-2}$. Then

$$V(\ell_1 \subset \Sigma) = \left[\begin{array}{cc|ccc} 0 & 0 & 0 & \cdots & 0 \\ \pm 1 & s & a_1 & \cdots & a_{2g} \\ \hline 0 & a_1 & & & \\ \vdots & \vdots & & W & \\ 0 & a_{2g} & & & \end{array}\right]$$

Because of Lemma 5.2, the Hoste formula (4) still holds. Therefore, the rest of the proof works out as in Proposition 5.1 to show that

$$\chi(I_*(M,P)) = -2s(\ell_1 \cup \ell_2 \subset \Sigma) = \frac{2 \cdot \lambda_L(M)}{|\operatorname{Tor}(H_1(M))|}$$

and that this is independent of the choice of admissible bundle which restricts to a trivial bundle on the exterior of $\ell_1 \cup \ell_2 \subset \Sigma$. □

Remark. The only remaining ambiguity in choosing an admissible bundle ad(P) over M has to do with the group $\operatorname{Ext}(H_1(M), \mathbb{Z}_2)$. If $H_1(M)$ has no 2-torsion, this group vanishes and Proposition 5.3 holds for an arbitrary admissible bundle.

6. CASE OF $b_1(M) = 3$

Let $\mathcal{L} = \ell_1 \cup \ell_2 \cup \ell_3$ be an algebraically split oriented three-component link in a rational homology sphere Σ such that each component of $\mathcal{L}$ is null-homologous. Let F_1, F_2, F_3 be Seifert surfaces of the knots ℓ_1, ℓ_2, ℓ_3, respectively, chosen so that $F_i \cap \ell_j = \emptyset$ for $i \neq j$. Define the Milnor triple linking number $\mu(\ell_1, \ell_2, \ell_3)$ as a signed count of points in the intersection $F_1 \cap F_2 \cap F_3$.

Given a closed oriented connected 3-manifold with $b_1(M) = 3$, by Lemma 2.1, there exists an algebraically split link $\mathcal{L} = \ell_1 \cup \ell_2 \cup \ell_3$ in a rational homology sphere Σ such that $M = \Sigma + 0 \cdot \ell_1 + 0 \cdot \ell_2 + 0 \cdot \ell_3$ and the components of $\mathcal{L}$ are all null-homologous. According to Lescop [9],

$$\lambda_L(M) = |\operatorname{Tor}(H_1(M))| \cdot (\mu(\ell_1, \ell_2, \ell_3))^2. \tag{6}$$

Again, $\lambda_L(M)$ is independent of the choice of surgery presentation as above.

Proposition 6.1. *Let* $M = \Sigma + 0 \cdot \ell_1 + 0 \cdot \ell_2 + 0 \cdot \ell_3$*, where* Σ *is a rational homology sphere,* $\ell_1, \ell_2, \ell_3 \subset \Sigma$ *are null-homologous knots such that* $\ell k_\Sigma(\ell_i, \ell_j)$ *=0 for* $i \neq j$*. Then*

$$\chi(I_*\,(M,P)) = -\frac{2 \cdot \lambda_L(M)}{|\operatorname{Tor}(H_1(M))|} \tag{7}$$

for any admissible bundle $\operatorname{ad}(P)$ *over* M *whose restriction to the exterior of* $\ell_1 \cup \ell_2 \cup \ell_3 \subset \Sigma$ *is trivial.*

Proof. Let P be a $U(2)$-bundle over M such that $w_2(\operatorname{ad}(P))$ evaluates nontrivially on all three homology classes obtained by capping off Seifert surfaces of ℓ_1, ℓ_2 and ℓ_3 by meridional disks of the surgery. By the Floer exact triangle,

$$\begin{aligned}\chi(I_*\,(M,P))) &= \chi(I_*\,((\Sigma + 0 \cdot \ell_1 + 0 \cdot \ell_2) - \ell_3)) - \chi(I_*\,(\Sigma + 0 \cdot \ell_1 + 0 \cdot \ell_2)) \\ &= -2s(\ell_1 \cup \ell_2 \subset (\Sigma - \ell_3)) + 2s(\ell_1 \cup \ell_2 \subset \Sigma) \\ &= -2 \cdot \ell k_{\Sigma - \ell_3}(c, c^+) + 2 \cdot \ell k_\Sigma(c, c^+),\end{aligned}$$

where c is the intersection circle $c = F_1 \cap F_2$ of the Seifert surfaces F_1 and F_2 chosen to intersect in a circle. We wish to identify the right hand side of this equation with $-2(\mu(\ell_1, \ell_2, \ell_3))^2$. By (5),

$$\ell k_{\Sigma - \ell_3}(c, c^+) - \ell k_\Sigma(c, c^+) = \ell k_\Sigma(c, \ell_3) \cdot \ell k_\Sigma(c^+, \ell_3),$$

therefore,

$$\chi(I_*\,(M,P))) = -2 \cdot \ell k_\Sigma(c, \ell_3) \cdot \ell k_\Sigma(c^+, \ell_3) = -2 \cdot \ell k(c, \ell_3)^2.$$

Since $\ell k_\Sigma(c, \ell_3)$= $c \cdot F_3 = \mu(\ell_1, \ell_2, \ell_3)$, the result follows. The independence of admissible bundle follows by the change of basis argument as before. $\square$

Remark. If $H_1(M)$ has no 2-torsion, by the same reasoning as in the remark after Proposition 5.1, the statement of Proposition 6.1 holds for any admissible bundle over M.

It is worth mentioning that Ruberman and Saveliev [12] showed that $1/2 \cdot \chi(I_*\,(M,P)) = \lambda_L(M) \mod 2$ for all M with $H_*(M) = H_*(T^3)$ using techniques different from ours.

7. CASE OF $b_1(M) \geq 4$

For all closed oriented connected 3-manifolds M with $b_1(M) = n \geq 4$, the Lescop invariant $\lambda_L(M)$ is known to vanish.

Proposition 7.1. *Let* $M = \Sigma + 0 \cdot \ell_1 + \ldots + 0 \cdot \ell_n$*, where* $n \geq 4$*,* Σ *is a rational homology sphere,* $\ell_1, \cdots, \ell_n \subset \Sigma$ *are null-homologous knots such that* $\ell k_\Sigma(\ell_i, \ell_j)$ *=0 for* $i \neq j$*. Then* $\chi(I_*(M, P)) = 0$ *for any admissible bundle* $\mathrm{ad}(P)$ *over* M *whose restriction to the exterior of* $\mathcal{L} = \ell_1 \cup \cdots \cup \ell_n$ *in* Σ *is trivial.*

Proof. Let P denote the $U(2)$-bundle over M such that $w_2(\mathrm{ad}(P))$ evaluates non-trivially on the homology classes obtained by capping off Seifert surfaces of the components ℓ_i, $i = 1, \ldots, n$, by meridional disks of the surgery. By the Floer exact triangle, if $n = 4$,

$$\begin{aligned}\chi(I_*(M,P)) &= \chi(I_*((\Sigma + 0\cdot\ell_1 + 0\cdot\ell_2 + 0\cdot\ell_3) - \ell_4)) \\ &\qquad - \chi(I_*(\Sigma + 0\cdot\ell_1 + 0\cdot\ell_2 + 0\cdot\ell_3)) \\ &\qquad\qquad = -2\cdot \ell k_{\Sigma-\ell_4}(c,\ell_3)^2 + 2\cdot \ell k_\Sigma(c,\ell_3)^2.\end{aligned}$$

The result now follows because

$$\ell k_{\Sigma-\ell_4}(c,\ell_3) - \ell k_\Sigma(c,\ell_3) = \ell k_\Sigma(c,\ell_4)\cdot \ell k_\Sigma(\ell_3,\ell_4) = 0,$$

using equation (5) and the fact that $\ell k(\ell_3, \ell_4) = 0$.

If $n \geq 5$, choose an admissible bundle $\mathrm{ad}(P)$ whose restriction to the exterior of $\mathcal{L} \subset \Sigma$ is trivial, and proceed by induction. Suppose that $\chi(I_*(M', P)) = 0$ for all M' with $b_1(M') = k$, where $4 \leq k \leq n-1$. Let $M = \Sigma + 0 \cdot \ell_1 + \ldots + 0 \cdot \ell_n$, where ℓ_i are null-homologous knots in a rational homology sphere Σ such that $\ell k_\Sigma(\ell_i, \ell_j)$ =0 for $i \neq j$. Then by the Floer exact triangle

$$\begin{aligned}\chi(I_*(M,P)) &= \chi(I_*((\Sigma + 0\cdot\ell_1 + \ldots + 0\cdot\ell_{n-1}) - \ell_n)) \\ &\qquad - \chi(I_*(\Sigma + 0\cdot\ell_1 + \ldots + 0\cdot\ell_{n-1})).\end{aligned}$$

Since

$$b_1((\Sigma - \ell_n) + 0\cdot\ell_1 + \ldots + 0\cdot\ell_{n-1}) = b_1(\Sigma + 0\cdot\ell_1 + \ldots + 0\cdot\ell_{n-1}) = n-1,$$

we conclude that $\chi(I_*(M,P)) = 0$. $\square$

Remark. As before, the result holds for all admissible bundles over M if $H_1(M)$ has no 2-torsion; the latter restriction can conjecturally be removed.

References

[1] S. Akbulut, J. McCarthy, *Casson's Invariant for Oriented Homology 3-Spheres: An Exposition.* Princeton Univ. Press, 1990.

[2] P. Braam, S. Donaldson, *Floer's work on instanton homology, knots, and surgery.* In: The Floer memorial volume, 195–256, Progr. Math. **133**, Birkhäuser, 1995.

[3] T. Cochran, *Derivatives of links: Milnors concordance invariants and Masseys products*, Mem. Amer. Math. Soc. **84** (1990), no. 427.

[4] S. Donaldson, *Floer homology groups in Yang-Mills theory.* Cambridge Univ. Press, 2002.

[5] A. Floer, *An instanton-invariant for 3-manifolds*, Comm. Math. Phys. **118** (1988), 215–240

[6] A. Floer, *Instanton homology, surgery and knots*, In: Geometry of low-dimensional manifolds, 1 (Durham, 1989). 97–114, London Math. Soc. Lecture Note Ser., **150**, Cambridge Uni. Press, 1990

[7] J. Hoste, *A formula for Cassons invariant*, Trans. Amer. Math. Soc. **297** (1986), 547–562

[8] P. Kronheimer and T. Mrowka, *Knots, sutures, and excision*, J. Differential Geom. **84** (2010), 301–364

[9] C. Lescop, *Global Surgery Formula for CassonWalker Invariant*, Annals of Mathematics Studies, Vol. 140. Princeton Univ. Press, 1996.

[10] M. Mariño and G. Moore, *3-manifold topology and the Donaldson-Witten partition function*, Nucl. Phys. **B547** (1999), 569–598

[11] K. Masataka, *Cassons knot invariant and gauge theory*, Topology Appl. **112** (2001), 111–135

[12] D. Ruberman and N. Saveliev, *Rohlins invariant and gauge theory. I. Homology 3-tori*, Comment. Math. Helv. **79** (2004), 618-646

[13] N. Saveliev, *Invariants for Homology 3-Spheres.* Springer, 2002.

[14] C. Taubes, *Casson's Invariant and Gauge Theory*, J. Differential Geom. **31** (1990), 547–599

Department of Mathematics
University of Miami
P.O.Box 249085
Coral Gables, FL 33124
E-mail address: poudelp@math.miami.edu